

ANTEPSIN-1000 Tablet

Sucralfate USP 1000 mg

Description:

ANTEPSIN is the brand name of Sucralfate, a basic Aluminium salt of Sucrose octasulphate. Sucralfate reacts with acid is converted into a sticky gel like substance (Sucralfate polyanion) at the site of ulcer and provides a proactive barrier against acid and pepsin of the stomach. Sucralfate is a locally acting drug.

Composition:

Each tablet contains Sucralfate USP 1000 mg.

Indication:

Gastric ulcer, duodenal ulcer and gastritis.

Dosage & Administration:

Adult: 1 tablet 4 times daily at least 1 hour before meal and at bed time. If necessary, the dose may be increased to a maximum of 8 tablets daily, in divided doses for 4-6 weeks or in resistant cases 12 weeks.

Elderly: No special dosage requirement is known but as with all medicines the lowest effective dose should be used.

Contra-indication:

There is no known contra-indication.

Precautions:

Antepsin should be given with caution in patient with renal disease.

Pregnancy and Lactation:

Sucralfate have not shown to cause adverse effects during pregnancy. Problems in human lactation have not been documented. However, risk/benefit must be considered when Sucralfate is administered to nursing mother.

Drug-interaction:

Concomitant use of Sucralfate with antacids may interfere with binding of Sucralfate to the gastric mucosa. Concomitant use of Antepsin may reduce the absorption of certain drugs such as phenytoin, tetracyclines and fat soluble vitamins.

Side-effects:

Side-effects with sucralfate are rare and mild. Rarely occurring side-effects are constipation, diarrhea, dizziness, drowsiness, dryness of mouth, indigestion, nausea, skin rash, stomach cramp or pain etc.

Packing:

Each box contains 10 blister strips of 10 tablets.

Manufactured by :



KEMIKO PHARMACEUTICALS LTD.

Bangladesh